

62735

(Pages : 3)

Name.....

60

Reg. No.....

SECOND SEMESTER B.C.A. DEGREE EXAMINATION, MAY 2014

(U.G.-CCSS)

Complementary Course

CA 2C 04—NUMERICAL METHODS IN C

Time : Three Hours

Maximum : 30 Weightage

Part A (Objective Type Questions)

Answer all twelve questions.

1. The numbers in the computer word can be stored in two forms. Which are they ?
2. Define the inherent error.
3. When we can say that ξ is a root of the equation $f(x) = 0$.
4. Define the central difference operator δ .
5. Write Newton's forward difference approximation of $0 (h^2)$.

6. What is the formula to find $\int_a^b f(x) dx$ using Simpson's rule ?

Fill in the blanks :

7. To avoid the difficulty of keeping every number less than 1 in magnitude during computation, most computers use _____ representation for a real number.
8. Bisection method is based on the repeated application of the _____ theorem.
9. In Gauss-Jordan elimination method the coefficient matrix is reduced to a _____ matrix.
10. If there are $n + 1$ distinct points $a \leq x_0 < x_1 < x_2 < \dots < x_n \leq b$, then the problem of Lagrange and Newton interpolation for the continuous function $f(x)$ on $[a, b]$ is to obtain $p(x)$ satisfying the conditions _____.
11. The Hermite interpolating polynomial interpolates not only the function $f(x)$ but also its _____ at a given set of tabular points.
12. The general problem of numerical integration is to find an approximate value of the integral $I = \int_a^b w(x) dx$ where $w(x) > 0$ in $[a, b]$.

(12 × ¼ = 3 weightage)

Turn over

Part B (Short Answer Questions)*Answer all nine questions.*

13. Find the decimal number corresponding to the binary number $(111.011)_2$.
14. Construct the difference table for the sequence of values $f(x) = (0, 0, 0, \varepsilon, 0, 0, 0)$.
15. Solve the equations $x + y = 2$ and $2x + 3y = 5$ by Gauss-Jordan method.
16. State intermediate value theorem.
17. Evaluate $\int_0^4 e^x dx$ by Simpson's '1/3' rule using the data $e = 2.72, e^2 = 7.39, e^3 = 20.09$
 $e^4 = 54.60$.
18. Perform 2 iterations of the bisection method to obtain a real root of the equation $x^3 - x - 11$
19. Solve $\frac{dy}{dx} = 1 - y, y(0) = 0$ using Euler's method. Find y at $x = 0.1$.
20. Find the n^{th} difference of e^x .
21. Show that $\mu = [1 + \delta^2 / 4]^{1/2}$.

 $(9 \times 1 = 9 \text{ weight})$ **Part C (Short Essay Questions)***Answer any five questions.*

22. Apply Cramer's rule to solve the equations, $3x + y + 2z = 3, 2x - 3y - z = 3$ and $x + 2y + z =$
23. Solve the following system of equations using Gaussian elimination method $x + y +$
 $2x - 3y + 4z = 13$ and $3x + 4y + 5z = 40$.
24. Construct Newton's forward interpolation polynomial for the following data :

$x :$	4	6	8	10
$y :$	1	3	8	16

25. Evaluate $\int_0^{10} \frac{dx}{1+x^2}$ by using Trapezoidal rule.

26. Using Taylor's method, find $y(0.1)$ from $\frac{dy}{dx} + 2xy = 1, y_0 = 0$.
27. Evaluate $\sqrt{12}$ to four places of decimals by Newton-Raphson method.
28. The equation $8x^3 - 12x^2 - 2x + 3 = 0$ has 3 real roots in the interval $[-2, 3]$. Find the intervals each of unit length containing each one of these roots.

(5 × 2 = 10 weightage)

Part D (Essay Questions)*Answer any two questions.*

29. (a) Write the Lagrange's interpolation formula.
- (b) Use Lagrange's formula to find the value of y at $x = 6$ from the following data :

x :	3	7	9	10
y :	168	120	72	63

30. (a) Find $y'(x)$ given :

x	:	0	1	2	3	4
$y(x)$:	1	1	15	40	85

- (b) The population of a certain town is shown in the following table :

Year x	:	1931	1941	1951	1961	1971
Population in 1961 y	:	40.62	60.80	79.95	103.56	132.65

31. (a) What is the relation between Runge-Kutta method and modified Euler's method.
- (b) Use Runge-Kutta method of the fourth order to find $y(0.1)$ given that :

$$\frac{dy}{dx} = \frac{1}{x+y}, y(0) = 1.$$

(2 × 4 = 8 weightage)